arXiv:0902.0086v1 [math.DG] 31 Jan 2009

Maurer–Cartan Forms of the Symmetry Pseudo-Group and the Covering of Plebañski's Second Heavenly Equation

Oleg I. Morozov

Department of Mathematics, Moscow State Technical University of Civil Aviation,
Kronshtadtskiy Blvd 20, Moscow 125993, Russia
oim@foxcub.org

Abstract. We derive Wahlquist–Estabrook forms of the covering of Plebañski's second heavenly equation from Maurer–Cartan forms of its symmetry pseudo-group.

AMS classification scheme numbers: 58H05, 58J70, 35A30

1. Introduction

In our preceding papers [16] – [19] it was shown that for a number of nonlinear partial differential equations (PDEs) with three independent variables Wahlquist–Estabrook forms of their coverings can be derived from Maurer–Cartan forms of their symmetry pseudo-groups. In this paper we consider Plebañski's second heavenly equation, [21],

$$u_{xz} = u_{ty} + u_{xx}\, u_{yy} - u_{xy}^2, \tag{1}$$

describing self-dual metrics in theory of gravitation. This equation can be obtained as the compatibility condition for the following system of PDEs, [8, 1], cf. [21, Eq. (3.13)]:

$$q_t = (u_{xy} - \lambda)\, q_x - u_{xx}\, q_y, \qquad q_z = u_{yy}\, q_x - (u_{xy} + \lambda)\, q_y, \tag{2}$$

where λ is an arbitrary constant. This condition is equivalent to the commutativity of the following four infinite-dimensional vector fields

$$\widetilde{D}_t = \bar{D}_t + \sum_{i,j \ge 0} \widetilde{D}_x^i \widetilde{D}_y^j \left((u_{xy} - \lambda)\, q_{1,0} - u_{xx}\, q_{0,1}\right) \frac{\partial}{\partial q_{i,j}},$$

$$\widetilde{D}_x = \bar{D}_x + \sum_{i,j \ge 0} q_{i+1,j} \frac{\partial}{\partial q_{i,j}}, \tag{3}$$

$$\widetilde{D}_y = \bar{D}_y + \sum_{i,j \ge 0} q_{i,j+1} \frac{\partial}{\partial q_{i,j}}, \tag{4}$$

$$\widetilde{D}_z = \bar{D}_z + \sum_{i,j \ge 0} \widetilde{D}_x^i \widetilde{D}_y^j \left(u_{yy}\, q_{1,0} - (u_{xy} + \lambda)\, q_{0,1}\right) \frac{\partial}{\partial q_{i,j}},$$

where $\bar{D}_t$, $\bar{D}_x$, $\bar{D}_y$ and $\bar{D}_z$ are restrictions of the total derivatives D_t, D_x, D_y and D_z to the infinite prolongation of (1). This construction is called a *covering*, [10] – [13]. Dually coverings can be defined by means of differential 1-forms called *Wahlquist–Estabrook forms*, [22]. For Eq. (1) an ideal of the Wahlquist–Estabrook forms is generated by the following forms:

$$\begin{aligned} \omega_{0,0} &= dq_{0,0} - \left((u_{xy} - \lambda)\, q_{1,0} - u_{xx}\, q_{0,1}\right) dt - q_{1,0}\, dx - q_{0,1}\, dy \\ &\qquad - \left(u_{yy}\, q_{1,0} - (u_{xy} + \lambda)\, q_{0,1}\right) dz, \\ \omega_{i,j} &= \widetilde{D}_x^i \widetilde{D}_y^j\, \omega_{0,0}, \qquad i, j \ge 0. \end{aligned} \tag{5}$$

In this work we establish that the form $\omega_{0,0}$ can be derived from Maurer–Cartan forms of the contact symmetry pseudo-group of Eq. (1).

2. Symmetry pseudo-groups of differential equations

Let $\pi : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R}^n$ be a vector bundle with the local base coordinates $(x^1, ..., x^n)$ and the local fibre coordinate u; then by $J^2(\pi)$ denote the bundle of the second-order jets of sections of π, with the local coordinates (x^i, u, u_i, u_{ij}), $i, j \in \{1, ..., n\}$. For every local section $(x^i, f(x))$ of π, denote by $j_2(f)$ the corresponding 2-jet $(x^i, f(x), \partial f(x)/\partial x^i, \partial^2 f(x)/\partial x^i \partial x^j)$. A differential 1-form ϑ on $J^2(\pi)$ is called a *contact* *form* if it is annihilated by all 2-jets of local sections: $j_2(f)^*\vartheta = 0$. In the local coordinates every contact 1-form is a linear combination of the forms $\vartheta_0 = du - u_i\,dx^i$, $\vartheta_i = du_i - u_{ij}\,dx^j$, $i, j \in \{1, ..., n\}$, $u_{ji} = u_{ij}$ (here and later we use the Einstein summation convention, so $u_i\,dx^i = \sum_{i=1}^{n} u_i\,dx^i$, etc.) A local diffeomorphism $\Delta : J^2(\pi) \to J^2(\pi)$, $\Delta : (x^i, u, u_i, u_{ij}) \mapsto (\bar{x}^i, \bar{u}, \bar{u}_i, \bar{u}_{ij})$, is called a *contact transformation* if for every contact 1-form $\bar{\vartheta}$ the form $\Delta^*\bar{\vartheta}$ is also contact. We denote by $\mathrm{Cont}(J^2(\pi))$ the pseudo-group of contact transformations on $J^2(\pi)$.

Let $\mathcal{H} \subset \mathbb{R}^{(2n+1)(n+3)(n+1)/3}$ be an open set with local coordinates a, b^i_k, c^i, f^{ik}, g_i, s_{ij}, w^k_{ij}, z_{ijk}, $i, j, k \in \{1, ..., n\}$, such that $a \neq 0$, $\det(b^i_k) \neq 0$, $f^{ik} = f^{ki}$, $z_{ijk} = z_{jik} = z_{ikj}$. Let (B^i_k) be the inverse matrix for the matrix (b^k_l), so $B^i_k\,b^k_l = \delta^i_l$. We consider the *lifted coframe*

$$\Theta_0 = a\,\vartheta_0, \quad \Theta_i = g_i\,\Theta_0 + a\,B^k_i\,\vartheta_k, \quad \Xi^i = c^i\,\Theta_0 + f^{ik}\,\Theta_k + b^i_k\,dx^k,$$

$$\Sigma_{ij} = s_{ij}\,\Theta_0 + w^k_{ij}\,\Theta_k + z_{ijk}\,\Xi^k + a\,B^k_i\,B^l_j\,du_{kl}, \tag{6}$$

$i \le j$, defined on $J^2(\pi) \times \mathcal{H}$. As it is shown in [15], the forms (6) are Maurer–Cartan forms for $\mathrm{Cont}(J^2(\pi))$, that is, a local diffeomorphism $\widehat{\Delta} : J^2(\pi) \times \mathcal{H} \to J^2(\pi) \times \mathcal{H}$ satisfies the conditions $\widehat{\Delta}^*\,\bar{\Theta}_0 = \Theta_0$, $\widehat{\Delta}^*\,\bar{\Theta}_i = \Theta_i$, $\widehat{\Delta}^*\,\bar{\Xi}^i = \Xi^i$, and $\widehat{\Delta}^*\,\bar{\Sigma}_{ij} = \Sigma_{ij}$ whenever it is projectable on $J^2(\pi)$, and its projection $\Delta : J^2(\pi) \to J^2(\pi)$ is a contact transformation.

The structure equations for $\mathrm{Cont}(J^2(\pi))$ read

$$\begin{aligned}
d\Theta_0 &= \Phi^0_0 \wedge \Theta_0 + \Xi^i \wedge \Theta_i,\\
d\Theta_i &= \Phi^0_i \wedge \Theta_0 + \Phi^k_i \wedge \Theta_k + \Xi^k \wedge \Sigma_{ik},\\
d\Xi^i &= \Phi^0_0 \wedge \Xi^i - \Phi^i_k \wedge \Xi^k + \Psi^{i0} \wedge \Theta_0 + \Psi^{ik} \wedge \Theta_k,\\
d\Sigma_{ij} &= \Phi^k_i \wedge \Sigma_{kj} - \Phi^0_0 \wedge \Sigma_{ij} + \Upsilon^0_{ij} \wedge \Theta_0 + \Upsilon^k_{ij} \wedge \Theta_k + \Lambda_{ijk} \wedge \Xi^k,
\end{aligned}$$

where the additional forms Φ^0_0, Φ^0_i, Φ^k_i, Ψ^{i0}, Ψ^{ij}, Υ^0_{ij}, Υ^k_{ij}, and Λ_{ijk} depend on differentials of the coordinates of $\mathcal{H}$.

Suppose $\mathcal{E}$ is a second-order differential equation in one dependent and n independent variables. We consider $\mathcal{E}$ as a sub-bundle in $J^2(\pi)$. Let $\mathrm{Cont}(\mathcal{E})$ be the group of contact symmetries for $\mathcal{E}$. It consists of all the contact transformations on $J^2(\pi)$ mapping $\mathcal{E}$ to itself. Let $\iota_0 : \mathcal{E} \to J^2(\pi)$ be an embedding and $\iota = \iota_0 \times \mathrm{id} : \mathcal{E} \times \mathcal{H} \to J^2(\pi) \times \mathcal{H}$. Maurer–Cartan forms of the pseudo-group $\mathrm{Cont}(\mathcal{E})$ can be obtained from the forms $\theta_0 = \iota^*\Theta_0$, $\theta_i = \iota^*\Theta_i$, $\xi^i = \iota^*\Xi^i$ and $\sigma_{ij} = \iota^*\Sigma_{ij}$ by means of Cartan's method of equivalence, [2]–[5], [7, 9, 20], see details and examples in [6, 16, 17, 19].

3. Symmetry pseudo-group and the covering of Plebañski's equation

Following the method outlined in the previous section we find the Maurer–Cartan forms and their structure equations for the symmetry pseudo-group of Eq. (1). The structure equations for the forms θ_0, θ_i, ξ^i, $i \in \{1, 2, 3\}$, read

$$d\theta_0 = \eta_5 \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3 + \xi^4 \wedge \theta_4,$$

$$
\begin{aligned}
d\theta_1 &= (\eta_5 - \eta_1) \wedge \theta_1 - \eta_3 \wedge \theta_4 - \eta_6 \wedge \theta_2 + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4 + 3\,\sigma_{22}) \wedge \theta_3 \\
&\quad + \xi^1 \wedge \sigma_{11} + \xi^2 \wedge \sigma_{12} + \xi^3 \wedge \sigma_{13} + \xi^4 \wedge \sigma_{14}, \\
d\theta_2 &= \tfrac{1}{3}\,(\eta_4 - 2\,\eta_1 + 2\,\eta_5) \wedge \theta_2 - \eta_3 \wedge \theta_3 + \xi^1 \wedge \sigma_{12} + \xi^2 \wedge \sigma_{22} + \xi^3 \wedge \sigma_{23} \\
&\quad + \xi^4 \wedge (\sigma_{13} + \sigma_{22} + \sigma_{33}), \\
d\theta_3 &= \tfrac{1}{3}\,(\eta_1 - 2\,\eta_4 + 2\,\eta_5) \wedge \theta_3 - \eta_2 \wedge \theta_2 + \xi^1 \wedge \sigma_{13} + \xi^2 \wedge \sigma_{23} + \xi^3 \wedge \sigma_{33} + \xi^4 \wedge \sigma_{34}, \\
d\theta_4 &= (\eta_5 - \eta_4) \wedge \theta_4 - \eta_2 \wedge \theta_1 + \tfrac{1}{3}\,(\eta_5 - 2\,\eta_1 + 4\,\eta_4 + 3\,\sigma_{33}) \wedge \theta_2 + \xi^1 \wedge \sigma_{14} \\
&\quad - (2\,\eta_2 + 2\,\eta_3 + \eta_6 - 2\,\sigma_{23}) \wedge \theta_3 + \xi^2 \wedge (\sigma_{13} + \sigma_{22} + \sigma_{33}) + \xi^3 \wedge \sigma_{34} + \xi^4 \wedge \sigma_{44}, \\
d\xi^1 &= \eta_1 \wedge \xi^1 + \eta_2 \wedge \xi^4, \\
d\xi^2 &= \eta_6 \wedge \xi^1 + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_1 - \eta_4) \wedge \xi^2 + \eta_2 \wedge \xi^3 + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_4 + 2\,\eta_1 + 3\,\sigma_{33}) \wedge \xi^4, \\
d\xi^3 &= \tfrac{1}{3}\,(4\,\eta_1 - 2\,\eta_4 - \eta_5 - 3\,\sigma_{22}) \wedge \xi^1 + \eta_3 \wedge \xi^2 + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_4 - \eta_1) \wedge \xi^3 \\
&\quad + (2\,\eta_2 + 2\,\eta_3 + \eta_6 - 2\,\sigma_{23}) \wedge \xi^4, \\
d\xi^4 &= \eta_3 \wedge \xi^1 + \eta_4 \wedge \xi^4.
\end{aligned}
$$

The involutive system of structure equations for this pseudo-group is given in Appendix. In the next calculations we use the following Maurer–Cartan forms only:

$$
\begin{aligned}
\xi^1 &= b_{11}\,dt + b_{14}\,dz, \\
\xi^2 &= v^{-1}\,\left(b_{11}\,dx + b_{14}\,dy - (b_{11}\,(w-1)\,u_{xy} + b_{14}\,u_{xx} + b_{41}\,v)\,dt\right. \\
&\qquad \left. -(b_{14}\,(w+1)\,u_{xy} - b_{11}\,u_{yy} + b_{44}\,v)\,dz\right), \\
\xi^3 &= v^{-1}\,\left(b_{41}\,dx + b_{44}\,dy + (b_{11}\,v - b_{41}\,(w-1)\,u_{xy} - b_{44}\,u_{xx})\,dt\right. \\
&\qquad \left. +(b_{14}\,v - b_{44}\,(w+1)\,u_{xy} + b_{41}\,u_{yy})\,dz\right), \\
\xi^4 &= b_{41}\,dt + b_{44}\,dz, \\
\eta_1 &= (b_{44}\,db_{11} - b_{41}\,db_{14})\,(b_{11}b_{44} - b_{14}b_{41})^{-1} + r_1\,\xi^1 + r_2\,\xi^4, \\
\eta_4 &= (b_{11}\,db_{44} - b_{14}\,db_{41})\,(b_{11}b_{44} - b_{14}b_{41})^{-1} - r_1\,\xi^1 - r_2\,\xi^4, \\
\eta_5 &= \ -3\,v^{-1}\,dv + \eta_1 + \eta_4,
\end{aligned}
$$

where b_{11}, b_{14}, b_{41}, b_{44}, v, w, r_1, r_2 are arbitrary parameters such that $b_{11}b_{44} - b_{14}b_{41} \neq 0$ and $v \neq 0$.

Direct computations prove the following

THEOREM. *Either substituting for* $v = q_{0,0}$, $b_{11} = q_{1,0}$, $b_{14} = q_{0,1}$, $w = \lambda\,u_{xy}^{-1}$ *into the linear combination*

$$\tfrac{1}{3}\,(\eta_1 + \eta_4 - \eta_5) - \xi^2 - \xi^4,$$

or substituting for $v = q_{0,0}$, $b_{41} = q_{1,0}$, $b_{44} = q_{0,1}$, $w = \lambda\,u_{xy}^{-1}$ *into the linear combination*

$$\tfrac{1}{3}\,(\eta_1 + \eta_4 - \eta_5) + \xi^1 - \xi^3$$

yields the form $q_{0,0}^{-1}\,\omega_{0,0}$ *proportional to the form (5), which is the generating form of the ideal of Wahquist–Estabrook forms of the covering (2) of Eq. (1).*

Another approach to computing Wahquist–Estabrook forms of coverings of PDEs from Maurer–Cartan forms of their symmetry pseudo-groups was proposed in [18]. We hope to apply this to Eq. (1) elsewhere.

References

[1] Bogdanov, L.V., Konopelchenko, B.G.: On the $\bar{\partial}$-dressing method applicaple to heavenly equation. Phys. Lett. A **345**, 137–143 (2005)

[2] Cartan, É.: Sur la structure des groupes infinis de transformations. Œuvres Complètes, Part II, **2**, 571–715. Gauthier - Villars, Paris (1953)

[3] Cartan, É.: Les sous-groupes des groupes continus de transformations. Œuvres Complètes, Part II, **2**, 719–856. Gauthier - Villars, Paris (1953)

[4] Cartan, É.: Les problèmes d'équivalence. Œuvres Complètes, Part II, **2**, 1311–1334. Gauthier - Villars, Paris (1953)

[5] Cartan, É.: La structure des groupes infinis. Œuvres Complètes, Part II, **2**, 1335–1384. Gauthier - Villars, Paris (1953)

[6] Fels, M., Olver, P.J.: Moving coframes. I. A practical algorithm. Acta. Appl. Math. **51**, 161–213 (1998)

[7] Gardner, R.B.: The Method of Equivalence and its Applications. CBMS–NSF regional conference series in applied math., SIAM, Philadelphia (1989)

[8] Husain, V.: Self-dual gravity and the chiral model. Phys. Rev. Lett., **72**, 800–803 (1994)

[9] Kamran, N.: Contributions to the Study of the Equivalence Problem of Élie Cartan and its Applications to Partial and Ordinary Differential Equations. Mem. Cl. Sci. Acad. Roy. Belg., **45**, Fac. 7 (1989)

[10] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal symmetries and the theory of coverings. Acta Appl. Math., **2**, 79–86 (1984)

[11] Krasil'shchik, I.S., Lychagin, V.V., Vinogradov, A.M.: Geometry of Jet Spaces and Nonlinear Partial Differential Equations. Gordon and Breach, New York (1986)

[12] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. Acta Appl. Math., **15**, 161–209 (1989)

[13] Krasil'shchik, I.S., Vinogradov, A.M. (eds.): Symmetries and Conservation Laws for Differential Equations of Mathematical Physics. Transl. Math. Monographs 182, Amer. Math. Soc., Providence (1999).

[14] Morozov, O.I.: Moving coframes and symmetries of differential equations. J. Phys. A, Math. Gen., **35**, 2965–2977 (2002)

[15] Morozov, O.I.: Contact-equivalence problem for linear hyperbolic equations. J. Math. Sci., **135**, 2680–2694 (2006)

[16] Morozov, O.I.: Coverings of differential equations and Cartan's structure theory of Lie pseudo-groups. Acta Appl. Math. **99**, 309–319 (2007)

[17] Morozov, O.I.: Cartan's structure theory of symmetry pseudo-groups, coverings and multi-valued solutions for the Khokhlov–Zabolotskaya equation, Acta Appl. Math. **101**, 231-241 (2008)

[18] Morozov, O.I.: Contact integrable extensions of symmetry pseudo-group and coverings of the r-th modified dispersionless Kadomtsev – Petviashvili equation Preprint `arXiv:0809.1218v1 [math.DG]` (2008)

[19] Morozov, O.I.: Cartan's structure of symmetry pseudo-group and coverings for the r-th modified dispersionless Kadomtsev–Petviashvili equation. Acta Appl. Math., accepted, 2009

[20] Olver, P.J.: Equivalence, Invariants, and Symmetry. Cambridge, Cambridge University Press (1995)

[21] Plebański, J.F.: Some solutions of complex Einstein equations, J. Math. Phys., **16**, 2395 – 2402 (1975)

[22] Wahlquist, H.D., Estabrook, F.B.: Prolongation structures of nonlinear evolution equations. J. Math. Phys., **16**, 1–7 (1975)

Appendix

The involutive system of structure equations for the symmetry pseudo-group of Eq. (1):

$$
\begin{aligned}
d\theta_0 &= \eta_5 \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3 + \xi^4 \wedge \theta_4,\\
d\theta_1 &= (\eta_5 - \eta_1) \wedge \theta_1 - \eta_3 \wedge \theta_4 - \eta_6 \wedge \theta_2 + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4 + 3\,\sigma_{22}) \wedge \theta_3\\
&\quad + \xi^1 \wedge \sigma_{11} + \xi^2 \wedge \sigma_{12} + \xi^3 \wedge \sigma_{13} + \xi^4 \wedge \sigma_{14},\\
d\theta_2 &= \tfrac{1}{3}\,(\eta_4 - 2\,\eta_1 + 2\,\eta_5) \wedge \theta_2 - \eta_3 \wedge \theta_3 + \xi^1 \wedge \sigma_{12} + \xi^2 \wedge \sigma_{22} + \xi^3 \wedge \sigma_{23}\\
&\quad + \xi^4 \wedge (\sigma_{13} + \sigma_{22} + \sigma_{33}),\\
d\theta_3 &= \tfrac{1}{3}\,(\eta_1 - 2\,\eta_4 + 2\,\eta_5) \wedge \theta_3 - \eta_2 \wedge \theta_2 + \xi^1 \wedge \sigma_{13} + \xi^2 \wedge \sigma_{23} + \xi^3 \wedge \sigma_{33} + \xi^4 \wedge \sigma_{34},\\
d\theta_4 &= (\eta_5 - \eta_4) \wedge \theta_4 - \eta_2 \wedge \theta_1 + \tfrac{1}{3}\,(\eta_5 - 2\,\eta_1 + 4\,\eta_4 + 3\,\sigma_{33}) \wedge \theta_2 + \xi^1 \wedge \sigma_{14}\\
&\quad - (2\,\eta_2 + 2\,\eta_3 + \eta_6 - 2\,\sigma_{23}) \wedge \theta_3 + \xi^2 \wedge (\sigma_{13} + \sigma_{22} + \sigma_{33}) + \xi^3 \wedge \sigma_{34} + \xi^4 \wedge \sigma_{44},\\
d\xi^1 &= \eta_1 \wedge \xi^1 + \eta_2 \wedge \xi^4,\\
d\xi^2 &= \eta_6 \wedge \xi^1 + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_1 - \eta_4) \wedge \xi^2 + \eta_2 \wedge \xi^3 + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_4 + 2\,\eta_1 + 3\,\sigma_{33}) \wedge \xi^4,\\
d\xi^3 &= \tfrac{1}{3}\,(4\,\eta_1 - 2\,\eta_4 - \eta_5 - 3\,\sigma_{22}) \wedge \xi^1 + \eta_3 \wedge \xi^2 + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_4 - \eta_1) \wedge \xi^3\\
&\quad + (2\,\eta_2 + 2\,\eta_3 + \eta_6 - 2\,\sigma_{23}) \wedge \xi^4,\\
d\xi^4 &= \eta_3 \wedge \xi^1 + \eta_4 \wedge \xi^4,\\
d\sigma_{11} &= \eta_7 \wedge \theta_1 + \eta_8 \wedge \theta_2 + \eta_9 \wedge \theta_3 + \eta_{10} \wedge \theta_4 + \eta_{11} \wedge \xi^1 + \eta_{12} \wedge \xi^2 + \eta_{13} \wedge \xi^3 + \eta_{14} \wedge \xi^4\\
&\quad + (\eta_5 - 2\,\eta_1) \wedge \sigma_{11} - 2\,\eta_6 \wedge \sigma_{12} + \tfrac{2}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4 + 3\,\sigma_{22}) \wedge \sigma_{13} - 2\,\eta_3 \wedge \sigma_{14},\\
d\sigma_{12} &= \eta_7 \wedge \theta_2 + \eta_{10} \wedge \theta_3 + \eta_{12} \wedge \xi^1 + (2\,\eta_7 - \eta_9) \wedge \xi^2 + \eta_{15} \wedge \xi^3 + \eta_{16} \wedge \xi^4 - (\eta_3 + \eta_6) \wedge \sigma_{22}\\
&\quad + \tfrac{1}{3}\,(2\,\eta_5 + \eta_4 - 5\,\eta_1) \wedge \sigma_{12} - 2\,\eta_3 \wedge \sigma_{13} + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_4 - 4\,\eta_1 + 3\,\sigma_{22}) \wedge \sigma_{23} - \eta_3 \wedge \sigma_{33},\\
d\sigma_{13} &= (\eta_{15} - \eta_8 - \eta_{10}) \wedge \theta_2 - \eta_7 \wedge \theta_3 + \eta_{13} \wedge \xi^1 + \eta_{15} \wedge \xi^2 + (\eta_{16} - 2\,\eta_7 + \eta_9 - \eta_{13}) \wedge \xi^3\\
&\quad + \eta_{17} \wedge \xi^4 - \eta_2 \wedge \sigma_{12} + \tfrac{2}{3}\,(\eta_5 - \eta_1 - \eta_4) \wedge \sigma_{13} - \eta_6 \wedge \sigma_{23} - \eta_3 \wedge \sigma_{34}\\
&\quad + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4 + 3\,\sigma_{22}) \wedge \sigma_{33},\\
d\sigma_{14} &= (\eta_{15} - \eta_8 - \eta_{10}) \wedge \theta_1 + (\eta_{16} + \eta_9 - \eta_{13}) \wedge \theta_2 - \eta_8 \wedge \theta_3 - \eta_7 \wedge \theta_4 + \eta_{14} \wedge \xi^1 + \eta_{16} \wedge \xi^2\\
&\quad + \eta_{17} \wedge \xi^3 + \eta_{18} \wedge \xi^4 - \eta_2 \wedge \sigma_{11} - \tfrac{1}{3}\,(\eta_5 - 2\,\eta_1 + 4\,\eta_4 - 3\,\sigma_{33}) \wedge \sigma_{12}\\
&\quad - 2\,(\eta_2 + \eta_3 + \eta_6 - \sigma_{23}) \wedge \sigma_{13} + (\eta_5 - \eta_1 - \eta_4) \wedge \sigma_{14} - \eta_6 \wedge (\sigma_{22} + \sigma_{33})\\
&\quad + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4 + 3\,\sigma_{22}) \wedge \sigma_{34} - \eta_3 \wedge \sigma_{44},\\
d\sigma_{22} &= (2\,\eta_7 - \eta_9) \wedge \xi^1 - \eta_{10} \wedge \xi^2 + \eta_7 \wedge \xi^3 + (2\,\eta_{15} - 2\,\eta_{10} - \eta_8) \wedge \xi^4 - 2\,\eta_3 \wedge \sigma_{23}\\
&\quad + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_1 + 2\,\eta_4) \wedge \sigma_{22},\\
d\sigma_{23} &= \eta_{15} \wedge \xi^1 + \eta_7 \wedge \xi^2 + (\eta_{15} - \eta_8 - \eta_{10}) \wedge \xi^3 + \eta_{19} \wedge \xi^4 - \eta_2 \wedge \sigma_{22} - \eta_3 \wedge \sigma_{33}\\
&\quad + \tfrac{1}{3}\,(\eta_5 - \eta_1 - \eta_4) \wedge \sigma_{23},\\
d\sigma_{33} &= (\eta_{16} - 2\,\eta_7 + \eta_9 - \eta_{13}) \wedge \xi^1 + (\eta_{15} - \eta_8 - \eta_{10}) \wedge \xi^2 + (\eta_{19} + \eta_7 - \eta_9 + \eta_{13} - \eta_{16}) \wedge \xi^3\\
&\quad + \eta_{20} \wedge \xi^4 - 2\,\eta_2 \wedge \sigma_{23} + \tfrac{1}{3}\,(\eta_5 + 2\,\eta_1 - 4\,\eta_4) \wedge \sigma_{33},\\
d\sigma_{34} &= (\eta_7 - \eta_9 + \eta_{13} - \eta_{16} + \eta_{19}) \wedge \theta_2 + (\eta_8 + \eta_{10} - \eta_{15}) \wedge \theta_3 + \eta_{17} \wedge \xi^1 + \eta_{19} \wedge \xi^2 + \eta_{20} \wedge \xi^3\\
&\quad + \eta_{21} \wedge \xi^4 - 2\,\eta_2 \wedge \sigma_{13} - \eta_2 \wedge \sigma_{22} - \tfrac{1}{3}\,(\eta_5 - 2\,\eta_1 + 4\,\eta_4) \wedge \sigma_{23}\\
&\quad - (3\,\eta_2 + 2\,\eta_3 + \eta_6 - 3\,\sigma_{23}) \wedge \sigma_{33} + \tfrac{1}{3}\,(2\,\eta_5 + \eta_1 - 5\,\eta_4) \wedge \sigma_{34},
\end{aligned}
$$

$$
\begin{aligned}
d\sigma_{44} &= (\eta_7 - \eta_9 + \eta_{13} - \eta_{16} + \eta_{19}) \wedge \theta_1 + (\eta_{20} - 2\,(\eta_8 + \eta_{10} - \eta_{15})) \wedge \theta_2 + \eta_{18} \wedge \xi^1 \\
&\quad - (\eta_9 - \eta_{13} + \eta_{16}) \wedge \theta_3 + (\eta_8 + \eta_{10} - \eta_{15}) \wedge \theta_4 + (\eta_{20} - \eta_8 - 2\,\eta_{10} + 2\,\eta_{15} + \eta_{17}) \wedge \xi^2 \\
&\quad + \eta_{21} \wedge \xi^3 + \eta_{22} \wedge \xi^4 - \tfrac{2}{3}\,(\eta_5 + 2\,\eta_1 - 4\,\eta_4 + 6\,\sigma_{33}) \wedge (\sigma_{13} + \sigma_{33}) - 2\,\eta_2 \wedge \sigma_{14} \\
&\quad - \tfrac{2}{3}\,(\eta_5 + 2\,\eta_1 - 4\,\eta_4 + 6\,\sigma_{33}) \wedge \sigma_{22} - 2\,(\eta_6 + 2\,(\eta_2 + \eta_3 - \sigma_{23})) \wedge \sigma_{34} \\
&\quad + (\eta_5 - 2\,\eta_4) \wedge \sigma_{44}, \\
d\eta_1 &= \eta_7 \wedge \xi^1 + (\eta_{15} - \eta_8 - \eta_{10}) \wedge \xi^4 + \eta_2 \wedge \eta_3, \\
d\eta_2 &= (\eta_{15} - \eta_8 - \eta_{10}) \wedge \xi^1 + (\eta_{19} + \eta_7 - \eta_9 + \eta_{13} - \eta_{16}) \wedge \xi^4 + (\eta_1 - \eta_4) \wedge \eta_2, \\
d\eta_3 &= \eta_{10} \wedge \xi^1 - \eta_7 \wedge \xi^4 + (\eta_4 - \eta_1) \wedge \eta_3, \\
d\eta_4 &= (\eta_8 + \eta_{10} - \eta_{15}) \wedge \xi^4 - \eta_7 \wedge \xi^1 - \eta_2 \wedge \eta_3, \\
d\eta_5 &= 0, \\
d\eta_6 &= \eta_8 \wedge \xi^1 + (\eta_7 - \eta_8 - \eta_{10} + \eta_{15}) \wedge \xi^3 + (\eta_9 - \eta_{13} + \eta_{16}) \wedge \xi^4 + \tfrac{1}{3}\,(\eta_6 - 4\,\eta_2 - 2\,\eta_3) \wedge \eta_1 \\
&\quad + \tfrac{1}{3}\,(2\,\eta_4 + \eta_5 + 3\,\sigma_{22}) \wedge \eta_2 + \tfrac{1}{3}\,(\eta_5 - 4\,\eta_4 + 3\,\sigma_{33}) \wedge \eta_3 - \tfrac{1}{3}(\eta_4 - \eta_5) \wedge \eta_6, \\
d\eta_7 &= \eta_{23} \wedge \xi^1 + \eta_{24} \wedge \xi^4 + \eta_7 \wedge \eta_1 - \eta_{10} \wedge \eta_2 + 2\,(\eta_{15} - \eta_8 - \eta_{10}) \wedge \eta_3, \\
d\eta_8 &= \eta_{25} \wedge \xi^1 + \eta_{23} \wedge \xi^2 + \eta_{24} \wedge \xi^3 + \eta_{26} \wedge \xi^4 - \tfrac{2}{3}\,(2\,\eta_8 + 5\,\eta_{10} - 4\,\eta_{15}) \wedge \eta_1 + \eta_2 \wedge \eta_9 \\
&\quad + 2\,(\eta_9 - \eta_{13} + \eta_{16}) \wedge \eta_3 + \tfrac{1}{3}\,(5\,\eta_8 + 8\,\eta_{10} - 4\,\eta_{15}) \wedge \eta_4 + \tfrac{1}{3}\,(\eta_8 + \eta_{10} - 2\,\eta_{15}) \wedge \eta_5 \\
&\quad - \eta_6 \wedge \eta_7 + 2\,(\eta_8 + \eta_{10} - \eta_{15}) \wedge \sigma_{22} - \eta_{10} \wedge \sigma_{33}, \\
d\eta_9 &= \eta_{27} \wedge \xi^1 + \eta_{28} \wedge \xi^2 - \eta_{23} \wedge \xi^3 - \eta_{25} \wedge \xi^4 - \tfrac{1}{3}\,(12\,\eta_7 + 7\,\eta_9) \wedge \eta_1 - (3\,\eta_8 + 2\,\eta_{10}) \wedge \eta_3 \\
&\quad + \eta_{10} \wedge (\eta_6 + 2\,\sigma_{23} - 2\,\eta_2) \\
&\quad + \tfrac{2}{3}\,(3\,\eta_7 - \eta_9) \wedge \eta_4 + \tfrac{1}{3}\,(3\,\eta_7 - \eta_9) \wedge \eta_5 + 3\,\eta_7 \wedge \sigma_{22}, \\
d\eta_{10} &= \eta_{28} \wedge \xi^1 - \eta_{23} \wedge \xi^4 + \eta_{10} \wedge (2\,\eta_1 - \eta_4) + 3\,\eta_3 \wedge \eta_7, \\
d\eta_{11} &= \eta_{23} \wedge \theta_1 + \eta_{25} \wedge \theta_2 + \eta_{27} \wedge \theta_3 + \eta_{28} \wedge \theta_4 + \eta_{29} \wedge \xi^1 + \eta_{30} \wedge \xi^2 + \eta_{31} \wedge \xi^3 + \eta_{32} \wedge \xi^4 \\
&\quad + (3\,\eta_{11} + 4\,\eta_{13}) \wedge \eta_1 + 3\,\eta_{14} \wedge \eta_3 - 2\,\eta_{13} \wedge \eta_4 - (\eta_{11} + \eta_{13}) \wedge \eta_5 \\
&\quad + 3\,(\eta_{12} \wedge (\eta_6 + \eta_8) - \eta_7 \wedge \sigma_{11} - \eta_9 \wedge \sigma_{13} - \eta_{10} \wedge \sigma_{14} - \eta_{13} \wedge \sigma_{22}), \\
d\eta_{12} &= \eta_{23} \wedge \theta_2 + \eta_{28} \wedge \theta_3 + \eta_{30} \wedge \xi^1 + (2\,\eta_{23} - \eta_{27}) \wedge \xi^2 + (\eta_{24} + \eta_{25} + \eta_{28}) \wedge \xi^3 \\
&\quad + (\eta_{26} - \eta_{27} + \eta_{31}) \wedge \xi^4 + \tfrac{8}{3}\,(\eta_{12} + \eta_{15}) \wedge \eta_1 + (\eta_{13} + 2\,\eta_{16}) \wedge \eta_3 - \tfrac{1}{3}\,(\eta_{12} + 4\,\eta_{15}) \wedge \eta_4 \\
&\quad - \tfrac{2}{3}\,(\eta_{12} + \eta_{15}) \wedge \eta_5 + 2\,(2\,\eta_7 - \eta_9) \wedge \eta_6 - 3\,\eta_7 \wedge \sigma_{12} - (\eta_8 + \eta_{10} + 2\,\eta_{15}) \wedge \sigma_{22} \\
&\quad - \eta_9 \wedge \sigma_{23} - \eta_{10} \wedge (3\,\sigma_{13} + \sigma_{33}), \\
d\eta_{13} &= \eta_{24} \wedge \theta_2 - \eta_{23} \wedge \theta_3 + \eta_{31} \wedge \xi^1 + (\eta_{24} + \eta_{25} + \eta_{28}) \wedge \xi^2 - (2\,\eta_{23} - \eta_{26}) \wedge \xi^3 + \eta_{33} \wedge \xi^4 \\
&\quad - \tfrac{1}{3}\,(16\,\eta_7 + 8\,\eta_9 - 3\,\eta_{13} + 8\,\eta_{16}) \wedge \eta_1 + \eta_{12} \wedge \eta_2 + 2\,\eta_{17} \wedge \eta_3 \\
&\quad + \tfrac{2}{3}\,(4\,\eta_7 - 2\,\eta_9 + 3\,\eta_{13} - 2\,\eta_{16}) \wedge \eta_4 + \tfrac{2}{3}\,(2\,\eta_7 - \eta_9 - \eta_{16}) \wedge \eta_5 + 2\,\eta_{15} \wedge (\eta_6 - \sigma_{12}) \\
&\quad + \eta_7 \wedge (\sigma_{13} + 4\,\sigma_{22}) + \eta_8 \wedge (2\,\sigma_{12} - \sigma_{23}) - \eta_9 \wedge (2\,\sigma_{22} + \sigma_{33}) + \eta_{10} \wedge (2\,\sigma_{12} - \sigma_{34}) \\
&\quad + 2\,(\eta_{13} - \eta_{16}) \wedge \sigma_{22}, \\
d\eta_{14} &= \eta_{24} \wedge \theta_1 + \eta_{26} \wedge \theta_2 - \eta_{25} \wedge \theta_3 - \eta_{23} \wedge \theta_4 + \eta_{32} \wedge \xi^1 + (\eta_{26} - \eta_{27} + \eta_{31}) \wedge \xi^2 + \eta_{33} \wedge \xi^3 \\
&\quad + \eta_{34} \wedge \xi^4 + \tfrac{2}{3}\,(\eta_{12} + 3\,\eta_{14} + 4\,\eta_{17}) \wedge \eta_1 + (\eta_{11} + 2\,\eta_{13}) \wedge \eta_2 + 2\,(\eta_{13} + \eta_{18}) \wedge \eta_3 \\
&\quad - \tfrac{1}{3}\,(4\,\eta_{12} - 3\,\eta_{14} + 4\,\eta_{17}) \wedge \eta_4 + \tfrac{1}{3}\,(\eta_{12} - 3\,\eta_{14} - 2\,\eta_{17}) \wedge \eta_5 + (\eta_{13} + 2\,\eta_{16}) \wedge \eta_6
\end{aligned}
$$

$$
\begin{aligned}
&+ \eta_7 \wedge \sigma_{14} + \eta_8 \wedge (2\,\sigma_{11} + \sigma_{13} - \sigma_{22} - \sigma_{33}) - \eta_9 \wedge (2\,\sigma_{12} + \sigma_{34}) + \eta_{10} \wedge (2\,\sigma_{11} - \sigma_{44}) \\
&+ \eta_{12} \wedge \sigma_{33} + 2\,((\eta_{13} + \eta_{16}) \wedge \sigma_{12} - \eta_{13} \wedge \sigma_{23} - \eta_{15} \wedge \sigma_{11} - \eta_{17} \wedge \sigma_{22}), \\
d\eta_{15} &= (\eta_{24} + \eta_{25} + \eta_{28}) \wedge \xi^1 + \eta_{23} \wedge \xi^2 + \eta_{24} \wedge \xi^3 + \eta_{35} \wedge \xi^4 - \tfrac{4}{3}\,(\eta_8 - \eta_{10} + 2\,\eta_{15}) \wedge \eta_1 \\
&+ (2\,\eta_7 - \eta_9) \wedge \eta_2 - (2\,\eta_7 - \eta_9 + \eta_{13} - \eta_{16} - \eta_{19}) \wedge \eta_3 + \tfrac{1}{3}\,(2\,(\eta_8 + \eta_{10}) - \eta_{15}) \wedge \eta_4 \\
&+ \tfrac{1}{3}\,(\eta_8 + \eta_{10} - 2\,\eta_{15}) \wedge \eta_5 - \eta_6 \wedge \eta_7 + 2\,(\eta_8 + \eta_{10} - \eta_{15}) \wedge \sigma_{22} - \eta_{10} \wedge \sigma_{33}, \\
d\eta_{16} &= \eta_{24} \wedge \theta_2 - \eta_{23} \wedge \theta_3 + (\eta_{26} - \eta_{27} + \eta_{31}) \wedge \xi^1 + (2\,\eta_{24} + \eta_{25}) \wedge \xi^2 + \eta_{35} \wedge \xi^3 + \eta_{36} \wedge \xi^4 \\
&+ \tfrac{1}{3}\,(4\,\eta_7 - 2\,\eta_9 + 5\,\eta_{16} + 4\,\eta_{19}) \wedge \eta_1 + (\eta_{12} + 2\,\eta_{15}) \wedge \eta_2 - (\sigma_{13} + 2\,\sigma_{33}) \wedge \eta_7 \\
&- (\eta_8 + 2\,\eta_{10} - 4\,\eta_{15} - 2\,\eta_{17} - \eta_{20}) \wedge \eta_3 - \tfrac{2}{3}\,(4\,\eta_7 - 2\,\eta_9 - \eta_{16} + \eta_{19}) \wedge \eta_4 \\
&+ \tfrac{1}{3}\,(2\,\eta_7 - \eta_9 - 2\,\eta_{16} - \eta_{19}) \wedge \eta_5 - (\eta_8 + 2\,\eta_{10} - 3\,\eta_{15}) \wedge \eta_6 - (2\,\sigma_{12} + \sigma_{23}) \wedge \eta_8 \\
&+ (\sigma_{22} + \sigma_{33}) \wedge \eta_9 - (2\,\sigma_{12} + \sigma_{34}) \wedge \eta_{10} + (\eta_{13} - \eta_{16} - \eta_{19}) \wedge \sigma_{22} - 2\,\eta_{15}(\wedge \sigma_{12} + \sigma_{23}), \\
d\eta_{17} &= (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \theta_2 - \eta_{24} \wedge \theta_3 + \eta_{33} \wedge \xi^1 + \eta_{35} \wedge \xi^2 - (2\,\eta_{24} + \eta_{25} + \eta_{33} - \eta_{36}) \wedge \xi^3 \\
&+ \eta_{37} \wedge \xi^4 + \tfrac{2}{3}\,(\eta_{15} + \eta_{17} + 2\,\eta_{20}) \wedge \eta_1 - (4\,\eta_7 - 2\,\eta_9 + \eta_{13} - 3\,\eta_{16}) \wedge \eta_2 \\
&- (4\,\eta_7 - 2\,\eta_9 + 2\,\eta_{13} - 2\,\eta_{16} - \eta_{21}) \wedge \eta_3 - \tfrac{1}{3}\,(4\,\eta_{15} - 5\,\eta_{17} + 2\,\eta_{20}) \wedge \eta_4 \\
&+ \tfrac{1}{3}\,(\eta_{15} - 2\,\eta_{17} - \eta_{20}) \wedge \eta_5 - (2\,\eta_7 - \eta_9 + \eta_{13} - \eta_{16} - \eta_{19}) \wedge \eta_6 - (\sigma_{12} + 3\,\sigma_{23}) \wedge \eta_9 \\
&+ (\sigma_{12} - 4\,\sigma_{23} - 2\,\sigma_{34}) \wedge \eta_7 - (\sigma_{13} - \sigma_{22} - 2\,\sigma_{33}) \wedge \eta_8 - (\sigma_{13} + \sigma_{22} + \sigma_{33}) \wedge \eta_{10} \\
&+ (\sigma_{12} - 3\,\sigma_{23}) \wedge \eta_{13} + (\sigma_{13} + \sigma_{22}) \wedge \eta_{15} - (\sigma_{12} + 3\,\sigma_{23}) \wedge \eta_{16} - \eta_{19} \wedge \sigma_{12} - \eta_{20} \wedge \sigma_{22}, \\
d\eta_{18} &= (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \theta_1 - (\eta_{25} + \eta_{33} - \eta_{36}) \wedge \theta_2 - \eta_{26} \wedge \theta_3 - \eta_{24} \wedge \theta_4 + \eta_{34} \wedge \xi^1 \\
&+ \eta_{36} \wedge \xi^2 + \eta_{37} \wedge \xi^3 + \eta_{38} \wedge \xi^4 + \tfrac{1}{3}\,(4\,\eta_{16} + 3\,\eta_{18} + 4\,\eta_{21}) \wedge \eta_1 + 2\,(\eta_{14} + 2\,\eta_{17}) \wedge \eta_2 \\
&+ (4\,\eta_{17} + \eta_{22}) \wedge \eta_3 - \tfrac{2}{3}\,(4\,\eta_{16} + 3\,\eta_{18} - \eta_{21}) \wedge \eta_4 + \tfrac{1}{3}\,(2\,\eta_{16} - 3\,\eta_{18} - \eta_{21}) \wedge \eta_5 \\
&- (\eta_8 + 2\,\eta_{10} - 2\,\eta_{15} - 3\,\eta_{17} - \eta_{20}) \wedge \eta_6 - \eta_7 \wedge (\sigma_{11} - 2\,\sigma_{44}) + \eta_8 \wedge (2\,\sigma_{12} + \sigma_{14} + 2\,\sigma_{34}) \\
&+ \eta_9 \wedge (\sigma_{11} - \sigma_{13} - 2\,\sigma_{22} - 2\,\sigma_{33}) + \eta_{10} \wedge (2\,\sigma_{12} + \sigma_{14}) - \eta_{15} \wedge (2\,\sigma_{12} + \sigma_{14}) \\
&- \eta_{13} \wedge (\sigma_{11} - \sigma_{13} - 2\,\sigma_{22} - 2\,\sigma_{33}) + \eta_{16} \wedge (\sigma_{11} - \sigma_{13} - 2\,\sigma_{22}) - 4\,\eta_{17} \wedge \sigma_{23} - \eta_{19} \wedge \sigma_{11} \\
&- \eta_{20} \wedge \sigma_{12} - \eta_{21} \wedge \sigma_{22}, \\
d\eta_{19} &= \eta_{35} \wedge \xi^1 + \eta_{24} \wedge \xi^2 + (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \xi^3 + \eta_{39} \wedge \xi^4 + \tfrac{1}{3}\,(2\,\eta_7 + \eta_{19}) \wedge \eta_1 \\
&- (3\,\eta_8 + 4\,\eta_{10} - 5\,\eta_{15}) \wedge \eta_2 - (2\,\eta_8 + 2\,\eta_{10} - 2\,\eta_{15} - \eta_{20}) \wedge \eta_3 - \tfrac{4}{3}\,(\eta_7 - \eta_{19}) \wedge \eta_4 \\
&+ \tfrac{1}{3}\,(\eta_7 - \eta_{19}) \wedge \eta_5 - (\eta_8 + \eta_{10} - \eta_{15}) \wedge \eta_6 - \eta_7 \wedge (\sigma_{22} - 2\,\sigma_{33}) \\
&+ 2\,(\eta_8 + \eta_{10} - \eta_{15}) \wedge \sigma_{23} + (\eta_9 - \eta_{13} + \eta_{16} - \eta_{19}) \wedge \sigma_{22}, \\
d\eta_{20} &= (\eta_{36} - 2\,\eta_{24} - \eta_{25} - \eta_{33}) \wedge \xi^1 + (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \xi^2 - (\eta_9 - \eta_{13} + \eta_{16} - 4\,\eta_{19}) \wedge \eta_2 \\
&+ (\eta_{24} + \eta_{25} + \eta_{33} - \eta_{36} + \eta_{39}) \wedge \xi^3 + \eta_{40} \wedge \xi^4 - \tfrac{2}{3}\,(\eta_8 + \eta_{10} - \eta_{15} + \eta_{20}) \wedge \eta_1 \\
&+ 2\,(\eta_7 - \eta_9 + \eta_{13} - \eta_{16} + \eta_{19}) \wedge \eta_3 + \tfrac{1}{3}\,(4\,(\eta_8 + \eta_{10} - \eta_{15}) + 7\,\eta_{20}) \wedge \eta_4 \\
&- \tfrac{1}{3}\,(\eta_8 + \eta_{10} - \eta_{15} + \eta_{20}) \wedge \eta_5 + (\eta_7 - \eta_9 + \eta_{13} - \eta_{16} + \eta_{19}) \wedge \eta_6 \\
&- 4\,(\eta_7 - \eta_9 + \eta_{13} - \eta_{16} + \eta_{19}) \wedge \sigma_{23} - 3\,(\eta_8 + \eta_{10} - \eta_{15}) \wedge \sigma_{33}, \\
d\eta_{21} &= (\eta_{24} + \eta_{25} + \eta_{33} - \eta_{36} + \eta_{39}) \wedge \theta_2 - (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \theta_3 + \eta_{37} \wedge \xi^1 + \eta_{39} \wedge \xi^2 \\
&+ \eta_{40} \wedge \xi^3 + \eta_{41} \wedge \xi^4 + \tfrac{1}{3}\,(4\,\eta_{19} - \eta_{21}) \wedge \eta_1 - (\eta_8 + 2\,\eta_{10} - 2\,\eta_{15} - 3\,\eta_{17} - 5\,\eta_{20}) \wedge \eta_2 \\
&- \tfrac{2}{3}\,(\eta_{19} - \eta_{21}) \wedge (4\,\eta_4 - \eta_5) - (\eta_7 - \eta_9) \wedge (3\,\sigma_{13} + 2\,\sigma_{22} + 2\,\sigma_{33}) + \eta_8 \wedge (2\,\sigma_{23} - 3\,\sigma_{34})
\end{aligned}
$$

$$
\begin{aligned}
&+ (\eta_{10} - \eta_{15}) \wedge (2\,\sigma_{23} - 3\,\sigma_{34}) - (\eta_{13} - \eta_{16}) \wedge (3\,\sigma_{13} + 2\,\sigma_{22} + 3\,\sigma_{33}) \\
&- \eta_{19} \wedge (3\,\sigma_{13} + 2\,\sigma_{22}) + \eta_{20} \wedge (4\,\eta_3 + 2\,\eta_6 - 5\,\sigma_{23}), \\
d\eta_{22} = {} & (\eta_{24} + \eta_{25} - \eta_{36} + \eta_{33} + \eta_{39}) \wedge \theta_1 + (2\,(\eta_{23} - \eta_{26} + \eta_{35}) + \eta_{40}) \wedge \theta_2 - 3\,(\eta_7 + \eta_{19}) \wedge \sigma_{14} \\
&+ (\eta_{25} + \eta_{33} - \eta_{36}) \wedge \theta_3 - (\eta_{23} - \eta_{26} + \eta_{35}) \wedge \theta_4 + \eta_{38} \wedge \xi^1 + (3\,\eta_{18} + 2\,\eta_{21}) \wedge \eta_2 \\
&+ (2\,\eta_{23} - \eta_{26} + 2\,\eta_{35} + \eta_{37} + \eta_{40}) \wedge \xi^2 + \eta_{41} \wedge \xi^3 + \eta_{42} \wedge \xi^4 + 3\,\eta_{21} \wedge (2\,\eta_3 + \eta_6 - 2\,\sigma_{23}) \\
&- 2\,(\eta_8 + 2\,(\eta_{10} - \eta_{15}) - \eta_{17} - \eta_{20}) \wedge \eta_1 + 3\,(\eta_9 - \eta_{13} + \eta_{16}) \wedge (\sigma_{14} + \sigma_{34}) \\
&+ (4\,(\eta_8 + 2\,(\eta_{10} - \eta_{15}) - \eta_{17} - \eta_{20}) + 3\,\eta_{22}) \wedge \eta_4 + 3\,\eta_{17} \wedge \sigma_{33} - 3\,\eta_{20} \wedge (\sigma_{13} + \sigma_{22}) \\
&- (\eta_8 + 2\,(\eta_{10} - \eta_{15}) - \eta_{17} - \eta_{20} + \eta_{22}) \wedge \eta_5 + 3\,\eta_8 \wedge (2\,(\sigma_{13} + \sigma_{22}) + \sigma_{33} - \sigma_{44}) \\
&+ 3\,(\eta_{10} - \eta_{15}) \wedge (2\,\sigma_{13} + 2\,\sigma_{22} - \sigma_{44}).
\end{aligned}
$$